

The exhaust valve of the North Atlantic

Agatha M. de Boer^{1,2}

and

Doron Nof^{1,3}

¹*Department of Oceanography 4320
The Florida State University
Tallahassee, Florida 32306-4320 U.S.A.
nof@ocean.fsu.edu*

²*Presently at: Atmospheric and Ocean Sciences Program
Princeton University/GFDL
P.O. Box 308, Forrestal Campus
Princeton, NJ 08540*

³*Also affiliated with The Geophysical Fluid Dynamics Institute, Florida State University*

Revised, September 2003

Abstract

During glacial periods, climate records are marked by large-amplitude oscillations believed to be a result of North Atlantic (NA) freshwater anomalies which weakened the thermohaline circulation (THC) and introduced instabilities. Such oscillations are absent from our present interglacial. With the aid of a semi-global analytical model, we propose that the Bering Strait (BS) acts like an exhaust valve for the above NA freshwater anomalies. Specifically, we suggest that large instabilities in the THC are only possible during glacial periods because, during these periods, the BS is closed. During interglacial periods (when the BS, the exhaust valve, is open), low salinity anomalies are quickly flushed out of the North Atlantic by the strong Southern Ocean winds.

1. Introduction. Recently, the difference between the climate stability during the Holocene and glacial periods (shown in **Fig. 1**) has been attributed to changes in the location of the North Atlantic Deep Water (NADW) formation region (e.g., Schmittner et al. 2002; Ganapolski and Rahmstorf 2001). It was also suggested that, during the glacial period, the meridional overturning cell was weak and, hence, less stable (e.g., Tziperman 1997). All of these results are based on numerical and climate models which display a very gradual glacial-interglacial transition. As such, they do not explain the extremely rapid climate transition so clearly visible in **Fig. 1** (left and right of the thick dashed line).

These relatively complex climate models also suffer from the familiar weakness that all general ocean circulation models suffer from – the results of simulations involving convection of roughly 10–20 Sv ($1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$) are often too sensitive to the specified vertical eddy diffusivities (roughly $0.5 \text{ cm}^2 \text{ s}^{-1}$) which are typically an order of magnitude higher than the observed values ($0.1 \text{ cm}^2 \text{ s}^{-1}$; see Bryan 1987). This is not a trivial matter considering that the commonly used numerical diffusivity of $0.5 \text{ cm}^2 \text{ s}^{-1}$ is associated with a conversion of as much as 10 Sv of upper ocean water to lower layer water or vice versa. (This scaling estimate is based on a thermocline 500 m deep and an Atlantic surface area of $5000 \text{ km} \times 20,000 \text{ km}$.) Even though attempts are being made to overcome these difficulties using non-uniform diffusivities, the issue is very troublesome particularly for

simulations involving flows through straits and passages. This is so because the transport in many passages is of the order of one Sverdrup, much smaller than the (excess) diffusion-induced transport. As a result, several numerical simulations predict flows (through straits) that are not even in the same direction as the observations (e.g., Nof and Van Gorder 2003).

Here, we present a simple analytical model related to the opening of the BS during the onset of the Holocene. We shall show that an open BS acts as an open exhaust valve allowing the flushing of anomalies out of the Atlantic. The analytics will be followed by process-oriented numerical simulations (using the primitive equations). We shall see that the model provides a compelling new explanation for the difference between the climate stability during glacials and interglacials and that it can explain the abrupt transition between the two regimes if (as argued below) the onset of the flow through the Bering Strait occurred suddenly. The change in stability will be related to our new Atlantic flushing mechanism whereas the sudden onset of the flow will be attributed to the initial jamming of the BS by icebergs and sea ice. Our model will clearly point to the physical processes responsible for the process – the strong southern winds (SW) and the geography of the American continent and the Atlantic Ocean.

2. The Bering Strait as an exhaust valve. Since the last glacial maximum (LGM), global sea level has risen by 130 m. The BS, being presently only 45 m deep, opened up to flow around 10.4 kyr BP (Before Present) which is coincident with the onset of the stable Holocene. We suggest that this opening took place abruptly (relative to the slow sea level rise) due to an initial jamming of the strait by icebergs and sea ice. We suppose that such a jamming resulted from the funnel-like shape of the surrounding continents. It persisted until the sea level difference between the Arctic and Pacific rose to a level high enough to break the “dam.” Such a scenario is not unusual. It has been argued that Lake Agassiz was dammed in a similar fashion and that icebergs also jammed the Denmark Straits (Van Kreveland et al. 2000). Unfortunately, because of the shallowness of the modern BS and

because of the currents associated with it, it is unlikely that any iceberg-generated scouring can still be identified on the bottom.

Ocean general circulation model (OGCM) studies indicate that NADW formation is reduced by the buoyancy provided by the BS throughflow (e.g., Wadley and Bigg 2002; Weijer et al. 2000; Reason and Power 1994). In line with this idea, Shaffer and Bendtsen (1994) employed a three-box model with a geostrophically controlled density driven BS flow to show that the overturning could be arrested by the freshwater flux. Here, a strikingly different mechanism, which considers the global ocean and the dynamical role of the open BS in the general oceanic circulation (rather than merely the buoyancy effect of an open BS) is proposed.

We consider the Americas to be a gigantic island (when the BS is open) and, as in Nof (2000, 2002), we apply an extension of the Island Rule (Godfrey 1989) which takes the NADW into account. The essence of the approach is that by (horizontally) integrating the (vertically integrated) equations of motion along a *closed* contour which does not pass through any frictional western boundary currents, the pressure term drops out and, as a result, one obtains a direct connection between the transports and the wind stress. Using the contour shown in **Fig. 2a**, we show that, in the absence of NADW formation, the strong winds in the southern boundary of the Atlantic Ocean would push about 4 Sv into the South Atlantic (SA) and then out of the North Atlantic (NA) into the Arctic and then the Pacific.

In view of this, any salinity anomalies in the NA which are strong enough to completely collapse the meridional overturning cell (MOC) would be quickly (within the horizontal advection time scale which is several years) flushed out of the Atlantic via the open BS so that the MOC will quickly recover (**Fig. 3**). On the other hand, when the BS is closed, the anomalies are trapped in the Atlantic and cannot be flushed out. The MOC can still recover (e.g., Schmittner et al. 2002; Bard 2002), but much more slowly than in the open BS case. Although not shown here (due to space limitations), small freshwater perturbations which are not sufficient to shut down the NADW completely are also quickly flushed out with an open BS (De Boer 2003). We argue that it is because of this

flushing that the perturbations to the temperature field are *large* prior to the opening of the BS and *small* after the opening (**Fig. 1**).

In our analytical model, we take not only the sea level to be continuous across the BS (**Fig. 2a**), but the entire vertically integrated pressure is also taken to be continuous (even though there can be a pressure exerted on the BS sill). This assumption can be relaxed without altering the fundamental results and our numerical simulations will show that the BS sill does not alter the results significantly. This, as well as other related issues, cannot be presented here in detail (due to space limitations) and, therefore, reference is made to de Boer and Nof (2003) where all of the needed clarifications are presented in detail. The southern flow into the Atlantic, Q_1 , differs from the Pacific inflow, Q_2 , by the amount of sinking within the Atlantic, W . Note that W is not a free parameter but rather is a function of the surface fluxes, the geography and the wind field. Following Nof (2000, 2002) we find that these are,

$$Q_1 = \frac{\oint \rho dr + \rho f_2 W}{\rho(f_2 - f_1)}, \quad Q_2 = \frac{\oint \rho dr + \rho f_1 W}{\rho(f_2 - f_1)}, \quad (1)$$

where $\oint \rho dr$ is the counterclockwise integrated wind stress along the path shown in **Fig. 2a**, ρ is the oceanic density and $f_{1,2}$ are the average Coriolis parameters at the southern and northern ends of the Atlantic, respectively (i.e., sections AB and CD). Both Q 's are positive towards the Atlantic and represent water from the surface to a depth above the topography. It is important to realize that (1) contains both wind and thermohaline processes.

Integration of 40 years of National Centers for Environmental Protection (NCEP) annual winds along the path shown in **Fig. 2a** gives 4 Sv for the first term in equations (1) (i.e., $\oint \rho dr / \rho(f_2 - f_1)$, where f_1 and f_2 are the Coriolis parameters at 45°S and 75°N, respectively). This 4 Sv is the amount that, in the absence of NADW (i.e., $W = 0$ and $Q_2 = -Q_1$), would be forced into the SA (Q_1) and out of the NA ($-Q_2$). This situation is shown in the right panel of **Fig. 2a**. Since the winds over the Arctic and NA are relatively weak, 60% of the 4 Sv is due to the strong SA winds, i.e., 60% of the wind stress contribution to

the integral comes from the integration along section AB (**Fig. 2**). (Note that we have made the plausible assumption here that the wind field has not changed much during the last 12,000 years.) With the present (nonzero) rate of NADW formation, Q_2 is almost 1 Sv (into the Atlantic) so that $W = 11$ Sv and $Q_1 = 10$ Sv. This situation is shown in the left panel of **Fig. 2a**. Note that, as discussed in Nof (2000), two thirds of the Agulhas rings contribution to the MOC is neglected here (because of our linearization). This amounts to a neglect of several Sverdrups in our calculation.

These considerations show that, in the absence of NADW formation (due to a strong freshwater anomaly), there is a reversal of the flow through the BS, i.e., without NADW, the flow is from the Arctic to the Pacific rather than the other way around. This is supported by our own process-oriented numerical simulations (**Fig. 3b**) employing the primitive equations without vertical eddy diffusivity. The “reduced gravity” model is only subject to wind forcing because we consider here the case of no NADW. The thin island represents the Americas and, as in the real world, the winds in the south are stronger than the winds in the north. Because of this asymmetry of the wind field, the flow around the island is counter-clockwise, implying a flow reversal through the BS (during NADW shut-off periods). This flow reversal was also noted in more complicated primitive equations models (Schiller et al. 1997). Furthermore, although there is no direct observational evidence for this flow reversal during a NADW shutdown, isotopic records of the northern Pacific strongly suggest that, in the early Holocene, North Pacific waters were freshened by flows from the Arctic Ocean (Gorbarenko 1996).

3. Discussion and summary. As mentioned, it is argued that our open BS flushing recovery time (several years) is much faster than the closed BS recovery time which is tens or hundreds of years (Stocker 2000; Schmittner et al. 2002; Van Kreveld et al. 2000). The latter is primarily due to diffusion (which gradually dissipates the freshwater anomaly), evaporation (which ultimately eliminates the anomaly), sea ice (which, through brine rejection, increases the salinity), strengthening of the local winds [which causes local upwelling which, in turn, increases the surface salinity reducing the density

difference that the cooling needs to overcome (Schiller et al. 1997)], and the cooling of the atmosphere due to a collapsed MOC and a reduction in the northward oceanic heat transport (Saravanan and McWilliams 1995). All of these recovery processes are considerably longer than our open valve rapid flushing mechanism.

In conclusion, an open BS acts like an open exhaust valve and allows the use of an integral constraint that links the flow through the BS to deep water formation in the NA. The interplay between the strength of the global winds and thermohaline-induced overturning act to stabilize (rather than destabilize) the climate.

4. Acknowledgments. We thank Andreas Thurnherr for helpful comments on the manuscript. This study was supported by the National Aeronautics and Space Administration under grants NAG5-10860 and NGT5-30164; National Science Foundation contracts OCE 9911324 and OCE-0241036; and Office of Naval Research grant N00014-01-0291.

5. References

- Bryan, F., 1987: Parameter sensitivity of primitive equation ocean general circulation models. *J. Phys. Oceanogr.*, **17**, 970-985.
- Bard, E., 2002: Climate shock: Abrupt changes over millennial time scales. *Physics Today*, **55**(12), 32-38.
- De Boer, A. M., 2003: The Bering Strait and the Southern Ocean winds' grip on the world climate. Ph.D. Dissertation, Florida State University, 87 pp.
- De Boer, A. M. and D. Nof, 2003: The Bering Strait's grip on the northern hemisphere climate. *Deep-Sea Res. I*, revised and resubmitted.
- Ganapolski, A. and S. Rahmstorf, 2001: Rapid changes of glacial climate simulated in a couple climate model. *Nature* **409**, 153-158.

- Godfrey, J. S., 1989: A Sverdrup model of the depth-integrated flow for the world ocean allowing for island circulations. *Geophys. Astrophys. Fluid Dynamics*, **45**, 89-112.
- Gorbarenko, S. A., 1996: Stable isotope and lithologic evidence of Late-Glacial and Holocene oceanography of the Northwestern Pacific and its marginal seas. *Quarter. Res.* **46**, 230-250.
- McManus, J. F., D. W. Oppo, and J. L. Cullen, 1999: A 0.5-million-year record of millennial-scale climate variability in the North Atlantic. *Science* **283**, 971-975.
- Nof, D., 2000: Does the wind control the import and export of the South Atlantic? *J. Phys. Oceanogr.* **30**(11), 2650-2667.
- Nof, D., 2002: Is there a meridional overturning cell in the Pacific and Indian Oceans? *J. Phys. Oceanogr.* **32**(6), 1947-1959.
- Nof, D. and S. Van Gorder, 2003: Did an open Panama Isthmus correspond to an invasion of Pacific water into the Atlantic? *J. Phys. Oceanogr.*, **33**(7), 1324-1336.
- Paul, A. and M. Schulz, 2002: Holocene climate variability on centennial-to-millennial time scales: 2. Internal and forced oscillations as possible causes. In *Climate Development and History of the North Atlantic Realm*. Springer-Verlag Berlin Heidelberg, 55-73.
- Reason, C. J. C and S. B. Power, 1994: The influence of the Bering Strait on the circulation in a coarse resolution global ocean model. *Climate Dyn.* **9**, 363-369.
- Saravanan, R. and J. C. McWilliams, 1995: Multiple equilibria, natural variability, and climate transitions in an idealized ocean-atmosphere model. *J. Climate* **8**, 2296-2323.

- Schiller, A., U. Mikolajewicz, R. and Voss, 1997: The stability of the North Atlantic thermohaline circulation in a coupled ocean-atmosphere general circulation model. *Climate Dyn.* **13**, 325-347.
- Schmittner, A., M. Yoshimori, and A. J. Weaver, 2002: Instability of glacial climate in a model of the ocean-atmosphere-cryosphere system. *Science* **295**, 1489-1493.
- Shaffer, G. and J. Bendtsen, 1994: Role of the Bering Strait in controlling North Atlantic ocean circulation and climate. *Nature* **367**, 354-357.
- Stocker, T. F. , 2000: Past and future reorganizations in the climate system. *Quatern. Sci. Rev.* **19**, 301-319.
- Tziperman, E., 1997: Inherently unstable climate behaviour due to weak thermohaline ocean circulation. *Nature* **386**, 592-595.
- Van Kreveld, S., M. Sarnthein, H. Erlenkeuser, P. Grootes, S. Jung, M. J. Nadeau, U. Pflaumann, and A. Voelker, 2000: Potential links between surging ice sheets, circulation changes, and the Dansgaard-Oeschger cycles in the Irminger Sea, 60-18 kyr. *Paleoceanography* **15**: 425-442.
- Wadley, M. R. and G. R. Bigg, 2002: Impact of flow through the Canacian Archipelago and Bering Strait on the North Atlantic and Arctic circulation. *Quart. J. Roy. Meteor. Soc.* **128**(585), 2187-2203.
- Weijer, W., W. P. M. de Ruijter, and H. A. Dijkstra, 2000: Stability of the Atlantic overturning circulation: Competition between Bering Strait freshwater flux and Agulhas heat and salt sources. *J. Phys. Oceanogr.* **31**(8), 2385-2402.






Fig. 1. Temperature record of Greenland and the North Atlantic [adapted from Bard (2002)]. Prior to the opening of the BS, large ice sheets and their induced albedo kept the mean temperature lower than it is today. The occurrence of high amplitude rapid oscillations decreased abruptly immediately after the opening of the BS, 10.4 kyr BP (shown with the thick dashed line). We argue that, although the meridional overturning cell (MOC) recovers from a freshwater-induced collapse in both the open and closed BS cases, the recovery in the open BS case is much quicker than that in the closed case. This is related to our proposed BS flushing mechanism which explains the very small fluctuations during the last 10,000 years compared to the large fluctuations in the period prior to that.

Fig. 2. (a) Left panel: The general flow pattern and the integration path around the Americas and the Atlantic (black dashed line) for the case of a convective Atlantic. $Q_{1,2}$ are the transports into the Atlantic from the south and north, respectively and W is the convection (indicated by $\square$). Right panel: The flow pattern in the no convection case ($W \equiv 0$).

(b) "Reduced gravity" numerical experiments showing how a freshwater anomaly (shaded region) which is strong enough to shut off the NADW is flushed out of the Atlantic within a short time. The island in the center represents the Americas and the northern gap represents the BS. The shown experiment does not involve a sill but experiments with a sill (protruding up to 50 m from the free surface) show very similar results with deviations of less than 40% from the no-sill values. The upper layer has one density and the anomaly is represented by dyed fluid. Even though the thermo-dynamics and the NADW shutoff are not explicitly present here, their absence has no bearing on the issue at hand because this single layer model accurately reflects relation (1) which does contain the thermodynamics. The wind profile reflecting the relatively weak (strong) winds in the northern (southern) Atlantic is shown in the upper left panel. The numerical parameters are the same as those given in Nof and Van Gorder (2003). The nondimensional time ($T/\varpi R_d$, where T is the dimensional time and R_d is the Rossby radius) is given in the left corner of each panel.






Fig. 3. The response of the MOC to a large NA freshwater anomaly in the open BS case (left) and the closed BS geometry (right). The colors blue, green, yellow to red represent increasing salinity (in that order). NADW formation is indicated with . Black arrows show the direction of flow and the blue double-headed arrows represent anomalous freshwater discharge. When the BS is open and the NADW is active, a mix of fresh Pacific and salty SA water sinks in the NA (I). A hypothetical freshwater anomaly (e.g., ice sheet discharge) renders the NA too fresh for deep convection (II). Our calculation implies that, under such conditions, the Southern Winds would flush the anomaly out of the NA and into the Pacific (III). Hence, the MOC recovers quickly [$O(10)$ years] and returns to the original steady state (I). On the other hand, when the BS is closed, a hypothetical freshwater anomaly (from an ice sheet) which freshens the NA to a degree that it is too fresh for deep convection (II) is trapped in the NA and cannot be flushed out (III). Under such conditions, the MOC can still recover but much more slowly (hundreds or thousands of years).